


# Investigating the Sources and Optical Importance of Brown Carbon during DC3 and SEAC4RS

Rodney J. Weber, Jiumeng Liu, Eric Scheuer, Jack Dibb, Luke Ziemba, Kenneth L. Thornhill, Bruce E. Anderson, Armin Wisthaler, Tomas Mikoviny, Joshua Schwarz, Ru-shan Gao, Tom Ryerson, Ilana Pollack, Jeff Peischl

## Objectives


- ❖ DC3: Assess Optical Importance of BrC in Background Continental Troposphere.
- ❖ SEAC4RS: Investigate BrC in biomass burning plumes:
  - ❖ Comparison of direct measured BrC to Ångström Exponents from optical instruments (PSAP and possibly PAS)
  - ❖ Prevalence of BrC relative to BC
  - ❖ Formation of BrC with plume aging (secondary BrC)?
  - ❖ BrC versus Mixing State (Shell Cores) and light absorption

## Method: Direct Measurement of BrC



- ❖ **Brown carbon determined from solution extracts**
- ❖ Isolate brown carbon from other light absorbers
- ❖ Highly wavelength-resolved


## DC3 Results Summary: No Plumes



See: Liu et al, Geophys. Res. Lett., 41, doi:10.1002/2013GL058976


## SEAC4RS RESULTS

### Use of 3-λ PSAP Data




Use of PSAP data averaged to filter sampling times to determine: AAE, b<sub>ap</sub> at 365 nm of BC, Total b<sub>ap</sub>


### Aug. 19 Flight: BrC and AAE



### BrC relative to BC vs Absorption Ångström Exponents




### Optical Closure: BrC & BC @ 365nm



Note on comparison between PAS (photo-acoustic instrument) and PSAP. Absorption coefficients are highly correlated (r > 0.95) and averaging over the PAS sampling period AAE<sub>PAS</sub>/AAE<sub>PSAP</sub> = 0.90

### Is BrC Formed as Smoke Plume Ages?



### Summary/Future Work

- BrC is a significant light absorbing component of biomass burning plumes
  - Good closure between light absorption from BrC and observed abs at 365 nm
  - ~70% of light absorption at 365 nm from BrC (lower % at higher wavelengths)
  - ~ ½ of BrC is water-soluble (Aug. 19 plumes).
- Possible evidence of shell-core effects on light absorption at high loadings?
- Secondary BrC production may occur in select plumes.
  - Will compare with evolution of other plume parameters to assess why.
- Future Work: Assess contribution of BrC in smoke plumes to:
  - Atmospheric stability (T profile), cloud microphysics, regional Climate via CMAQ